

Indian Institute of Technology Roorkee
MAI-101(Mathematics-1)
Autumn Semester: 2024–25
Assignment-9: Vector Calculus I (Gradient, Divergence, Curl etc.)

Notation: $\mathbf{i} = \vec{i}$, $\mathbf{j} = \vec{j}$ and $\mathbf{k} = \vec{k}$ are the unit vectors along x , y and z -axes, respectively. Boldface letters represent vectors.

1. Show that

- (i) the necessary and sufficient condition for the vector function $\mathbf{u}(t) = u_1(t)\mathbf{i} + u_2(t)\mathbf{j} + u_3(t)\mathbf{k}$ to be a constant is that $\frac{d\mathbf{u}}{dt} = \mathbf{0}$.
- (ii) the necessary and sufficient condition for the vector function $\mathbf{u}(t) = u_1(t)\mathbf{i} + u_2(t)\mathbf{j} + u_3(t)\mathbf{k}$ to have constant magnitude is that $\mathbf{u} \cdot \frac{d\mathbf{u}}{dt} = 0$.
- (iii) the necessary and sufficient condition for the vector function $\mathbf{u}(t) = u_1(t)\mathbf{i} + u_2(t)\mathbf{j} + u_3(t)\mathbf{k}$ to have constant direction is $\mathbf{u} \times \frac{d\mathbf{u}}{dt} = \mathbf{0}$.

2. (i) If $\mathbf{r} = \mathbf{a}e^{nt} + \mathbf{b}e^{-nt}$, where \mathbf{a} and \mathbf{b} are constant vectors, show that $\frac{d^2\mathbf{r}}{dt^2} = n^2\mathbf{r}$.

(ii) If $\mathbf{r} = (\cos nt)\mathbf{i} + (\sin nt)\mathbf{j}$, show that $\mathbf{r} \times \frac{d\mathbf{r}}{dt} = n\mathbf{k}$.

3. Let $[\mathbf{a} \ \mathbf{b} \ \mathbf{c}] = \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$, for $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$, $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$

and $\mathbf{c} = c_1\mathbf{i} + c_2\mathbf{j} + c_3\mathbf{k}$. Given $\mathbf{r} = a \cos t\mathbf{i} + a \sin t\mathbf{j} + bt\mathbf{k}$, show that

- (i) $\mathbf{r}^2 = \mathbf{r} \cdot \mathbf{r} = a^2 + b^2t^2$,
- (ii) $|\mathbf{r}' \times \mathbf{r}''|^2 = a^2(a^2 + b^2)$,
- (iii) $[\mathbf{r}' \ \mathbf{r}'' \ \mathbf{r}'''] = a^2b$,

where $\mathbf{r}' = \frac{d\mathbf{r}}{dt}$, $\mathbf{r}'' = \frac{d^2\mathbf{r}}{dt^2}$ and $\mathbf{r}''' = \frac{d^3\mathbf{r}}{dt^3}$.

4. (i) If $\varphi = 2xz^4 - x^2y$, find $\nabla\varphi$ and $|\nabla\varphi|$ at the point $(2, -2, 1)$.

(ii) If $\nabla\varphi = (y + y^2 + z^2)\mathbf{i} + (x + z + 2xy)\mathbf{j} + (y + 2zx)\mathbf{k}$, find φ such that $\varphi(1, 1, 1) = 3$.

(iii) If $\varphi = (x^2 + y^2 + z^2)e^{-\sqrt{x^2+y^2+z^2}}$, find $\nabla\varphi$.

5. If $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and $|\mathbf{r}| = r$, then show that

- (i) $\nabla r^n = nr^{n-2}\mathbf{r}$,
- (ii) $\nabla \left(\frac{1}{r} \right) = -\frac{\mathbf{r}}{r^3}$,

$$(iii) \nabla f(r) = \frac{f'(r)}{r} \mathbf{r}, \quad \nabla f(r) \times \mathbf{r} = \mathbf{0},$$

$$(iv) \nabla[\mathbf{r} \cdot \mathbf{a} \cdot \mathbf{b}] = \mathbf{a} \times \mathbf{b},$$

where \mathbf{a} and \mathbf{b} are constant vectors.

6. (i) Find the directional derivative of $\varphi = x^2 - 2y^2 + 4z^2$ at $(1, 1, -1)$ in the direction of $2\mathbf{i} + 2\mathbf{j} + \mathbf{k}$.
- (ii) Find the directional derivative of $\varphi = x^2(y + z)$ at $(1, 1, 0)$ in the direction of the line joining the origin to the point $(2, -1, 2)$.
- (iii) Find the directional derivative of the function $\varphi = x^2 - y^2 + 2z^2$ at the point $P(1, 2, 3)$ in the direction of the line PQ , where Q is the point $(5, 0, 4)$.
- (iv) Find the direction along which the directional derivative of the function $\varphi = xy + 2yz + 3xz$ is greatest at the point $(1, 1, 1)$. Also find the greatest directional derivative.
- (v) Find the directional derivative of $\varphi = 4xz^3 - 3xyz^2$ at $(2, -1, 1)$, along z -axis.
7. Prove that the gradient is a vector perpendicular to the level surface $\varphi(x, y, z) = c$, here c is a constant.
8. (i) Find the unit vector normal to the level surface $xy + y^2 - z^2 = 5$ at $(1, 2, 1)$.
- (ii) Find the angle between the surfaces $x^2 + y^2 + z^2 = 9$ and $z = x^2 + y^2 - 3$ at the point $(2, -1, 2)$.
9. If \mathbf{a} is a constant vector and $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ with $r = |\mathbf{r}|$, then show that
 - (i) $\text{div}(\mathbf{r} \times \mathbf{a}) = 0$, i.e., $\mathbf{r} \times \mathbf{a}$ is solenoidal,
 - (ii) $\text{curl}(\mathbf{r} \times \mathbf{a}) = -2\mathbf{a}$ or $\nabla \times (\mathbf{a} \times \mathbf{r}) = 2\mathbf{a}$,
 - (iii) $\text{grad}(\mathbf{a} \cdot \mathbf{r}) = \mathbf{a}$,
 - (iv) $\nabla \cdot (r^2 \mathbf{a}) = 2\mathbf{a} \cdot \mathbf{r}$.
10. (i) Determine a so that the vector $\mathbf{F} = (z + 3y)\mathbf{i} + (x - 2z)\mathbf{j} + (x + az)\mathbf{k}$ is solenoidal.
- (ii) Find the value of a if $\mathbf{F} = (axy - z^2)\mathbf{i} + (x^2 + 2yz)\mathbf{j} + (y^2 - axz)\mathbf{k}$ is irrotational.
- (iii) A field \mathbf{F} is of the form $\mathbf{F} = (6xy + z^3)\mathbf{i} + (3x^2 - z)\mathbf{j} + (3xz^2 - y)\mathbf{k}$. Show that \mathbf{F} is a conservative field (i.e., \mathbf{F} is irrotational) and find its scalar potential.
11. If \mathbf{F} is a differentiable vector function and φ is a differentiable scalar function, then prove that
 - (i) $\text{div}(\varphi \mathbf{F}) = \text{grad } \varphi \cdot \mathbf{F} + \varphi \text{div } \mathbf{F}$ or $\nabla \cdot (\varphi \mathbf{F}) = \nabla \varphi \cdot \mathbf{F} + \varphi \nabla \cdot \mathbf{F}$,
 - (ii) $\text{curl}(\varphi \mathbf{F}) = \varphi \text{curl } \mathbf{F} + \text{grad } \varphi \times \mathbf{F}$ or $\nabla \times (\varphi \mathbf{F}) = \varphi (\nabla \times \mathbf{F}) + (\nabla \varphi) \times \mathbf{F}$.
12. For $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, show that
 - (i) $\nabla \cdot \left(\frac{\mathbf{r}}{r^3} \right) = 0$,

(ii) $\nabla \cdot (r^3 \mathbf{r}) = 6r^3$,

where $r = |\mathbf{r}|$ and \mathbf{a} is a constant vector.

13. If \mathbf{a} is a constant vector, then prove that

$$\text{curl} \left(\frac{\mathbf{a} \times \mathbf{r}}{r^3} \right) = -\frac{\mathbf{a}}{r^3} + \frac{3\mathbf{r}}{r^5}(\mathbf{a} \cdot \mathbf{r}).$$

14. If \mathbf{F} is a vector function, prove that $\text{curl}(\text{curl } \mathbf{F}) = \text{grad}(\text{div } \mathbf{F}) - \nabla^2 \mathbf{F}$, where $\nabla^2 = \nabla \cdot \nabla$.

15. If $\mathbf{F} = 2yz\mathbf{i} + x^2y\mathbf{j} + xz^2\mathbf{k}$, $\mathbf{G} = x^2\mathbf{i} + yz\mathbf{j} + xy\mathbf{k}$, and $\varphi = 2x^2yz^3$, find

(i) $(\mathbf{F} \cdot \nabla)\varphi$ (ii) $(\mathbf{F} \times \nabla)\varphi$ (iii) $(\nabla \times \mathbf{F}) \times \mathbf{G}$.

Answers.

4. (i) $\nabla\varphi|_{(2,-2,1)} = 10\mathbf{i} - 4\mathbf{j} + 16\mathbf{k}$, $|\nabla\varphi| = 2\sqrt{93}$. (ii) $\varphi = xy + xy^2 + xz^2 + yz - 1$.
 (iii) $e^{-\sqrt{x^2+y^2+z^2}}(2 - \sqrt{x^2+y^2+z^2})(x\mathbf{i} + y\mathbf{j} + z\mathbf{k})$.

6. (i) -4 . (ii) $\frac{5}{3}$. (iii) $\frac{28}{\sqrt{21}}$. (iv) $4\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}$, $5\sqrt{2}$. (v) 36 .

8. (i) $\frac{2\mathbf{i}+5\mathbf{j}-2\mathbf{k}}{\sqrt{33}}$. (ii) $\theta = \cos^{-1} \left(\frac{8}{3\sqrt{21}} \right)$.

10. (i) $a = 0$. (ii) $a = 2$. (iii) $\varphi = 3x^2y + xz^3 - yz + C$.

15. (i) $8xy^2z^4 + 2x^4yz^3 + 6x^3yz^4$.
 (ii) $[6x^4y^2z^4 - 2x^3z^5]\mathbf{i} - [12x^2y^2z^3 - 4x^2yz^5]\mathbf{j} + [2x^2yz^4 - 4x^3y^2z^3]\mathbf{k}$.
 (iii) $[2xy^2 - xyz^2 - 2xy^2z + 2yz^2]\mathbf{i} - [2x^3y - 2x^2z]\mathbf{j} + [x^2z^2 - 2x^2y]\mathbf{k}$.